

## Publications

1. Hochhaus A, Weisser A, La Rosee P, Emig M, Müller MC, Saussele S, Reiter A, Kuhn C, Berger U, Hehlmann R, Cross NC.  
Detection and quantification of residual disease in chronic myelogenous leukemia.  
Leukemia 2000; 14(6):998-1005.
2. Saussele S, Weisser A, Müller MC, Emig M, La Rosee P, Paschka P, Kuhn C, Willer A, Hehlmann R, Hochhaus A.  
Frequent polymorphism in BCR exon b2 identified in BCR-ABL positive and negative individuals using fluorescent hybridization probes.  
Leukemia 2000 Nov;14(11):2006-10
3. de la Fuente J, Merx K, Steer EJ, Müller MC, Szydlo RM, Maywald O, Berger U, Hehlmann R, Goldman JM, Cross NC, Melo JV, Hochhaus A.  
ABL-BCR expression does not correlate with deletions on the derivative chromosome 9 or survival in chronic myeloid leukemia.  
Blood 2001 Nov 1;98(9):2879-80.
4. Merx K, Müller MC, Kreil S, Lahaye L, Paschka P, Schoch C, Weisser A, Kuhn C, Berger U, Gschaidmeier H, Hehlmann R, and Hochhaus A.  
Early Reduction of BCR-ABL mRNA Transcript Levels Predicts Cytogenetic Response to Imatinib in Chronic Phase CML Patients.  
Leukemia 2002 Sep;16(9):1579-83.
5. Hochhaus A, Berger U, Reiter A, Engelich G, Lahaye T, Kreil S, Merx K, Müller MC, Hehlmann R.  
Aktuelle Therapiekonzepte bei chronischer myeloischer Leukämie.  
Internist 2002; 43:1228-1244.
6. Müller MC, Merx K, Weißer A, Kreil S, Lahaye T, Hehlmann R, and Hochhaus A.  
Improvement of Molecular Monitoring of Residual Disease in Leukemias by Bedside RNA Stabilization.  
Leukemia 2002 Dec; 16(12): 2395-2399.
7. Müller MC, Lahaye T, Hochhaus A.  
Resistenz auf tumorspezifische Therapie mit Imatinib durch klonale Selektion mutierter Zellen.  
Dtsch Med Wochenschr. 2002 Oct 18;127(42):2205-7.
8. Hochhaus A, Müller MC, Hehlmann R.

Diagnostik und Therapie der chronischen myeloischen Leukämie.  
Onkologie 2002; 8(7):693-707.

9. Hochhaus A, Kreil S, Corbin AS, La Rosée P, Müller MC, Lahaye T, Hanfstein B, Schoch C, Cross NCP, Berger U, Gschaidmeier H, Druker BJ and Hehlmann R.  
Molecular and chromosomal mechanisms of resistance to imatinib (STI571) therapy.  
Leukemia 2002 Nov; 16(11): 2190-2196.
10. Burchert A, Woelfl S, Schmidt M, Brendel C, Dennecke B, Cai D, Odynova L, Lahaye T, Müller MC, Berg T, Gschaidmeier H, Wittig B, Hehlmann R, Hochhaus A, and Neubauer A.  
Interferon-alfa, but not the ABL-kinase inhibitor imatinib (STI571), induces expression of myeloblastin and a specific T-cell response in chronic myeloid leukemia.  
Blood 2003;101 259-264 (Blood 2002 Jun 28 (epub ahead of print)).
11. Sacha T, Hochhaus A, Hanfstein B, Müller MC, Rudzki Z, Czopek J, Wolska-Smolen T, Czekalska S, Salamanchuk Z, Jakobczyk M, and Skotnicki AB.  
ABL-kinase domain point mutation as a cause of imatinib (STI571) resistance in CML patient who progress to myeloid blast crisis.  
Leukemia Research 2003 Dec; 27(12): 1163-1166.
12. Paschka P, Müller MC, Merx K, Kreil S, Schoch C, Lahaye T, Weisser A, Petzold A, König H, Berger U, Gschaidmeier H, Hehlmann R and Hochhaus A.  
Molecular monitoring of response to imatinib (Glivec®) in CML patients pretreated with interferon alpha. Low levels of residual disease are associated with continuous remission.  
Leukemia 2003 Sep; 17(9): 1687-1694.
13. Müller MC, Gattermann N, Lahaye T, Deininger MW, Berndt A, Fruehauf S, Neubauer A, Fischer T, Hossfeld DK, Schneller F, Krause SW, Nerl C, Sayer HG, Ottmann OG, Waller C, Aulitzky W, le Coutre P, Freund M, Merx K, Paschka P, König H, Kreil S, Berger U, Gschaidmeier H, Hehlmann R, Hochhaus A.  
Dynamics of BCR-ABL mRNA expression in first-line therapy of chronic myelogenous leukemia patients with imatinib or interferon alpha/ara-C.  
Leukemia 2003 Dec; 17(12): 2392-2400.
14. Müller MC, Hördt T, Paschka P, Merx K, La Rosée P, Hehlmann R, Hochhaus A.  
Standardization of preanalytical factors for minimal residual disease analysis in chronic myelogenous leukemia.  
Acta haematol. 2004; 112(1-2):30-3.
15. Lahaye T, Riehm B, Berger U, Paschka P, Müller MC, Kreil S, Merx K, Schwindel U, Schoch C, Hehlmann R, Hochhaus A.

Response and resistance in 300 patients with BCR-ABL-positive leukemias treated with imatinib in a single center: a 4.5-year follow-up.  
Cancer 2005 Apr 15;103(8):1659-69.

16. Frank O, Brors B, Fabarius A, Li L, Haak M, Merk S, Schwindel U, Zheng C, Müller MC, Gretz N, Hehlmann R, Hochhaus A, Seifarth W.  
Gene expression signature of primary imatinib-resistant chronic myeloid leukemia patients.  
Leukemia. 2006 Aug;20(8):1400-7. Epub 2006 May 25.
17. Roeder I, Horn M, Glauche I, Hochhaus A, Mueller MC, Loeffler M.  
Dynamic modeling of imatinib-treated chronic myeloid leukemia: functional insights and clinical implications.  
Nat Med. 2006 Oct;12(10):1181-4. Epub 2006 Oct 1.
18. Walz C, Curtis C, Schnittger S, Schultheis B, Metzgeroth G, Schoch C, Lengfelder E, Erben P, Müller MC, Haferlach T, Hochhaus A, Hehlmann R, Cross NC, Reiter A.  
Transient response to imatinib in a chronic eosinophilic leukemia associated with ins(9;4)(q33;q12q25) and a CDK5RAP2-PDGFR A fusion gene.  
Genes Chromosomes Cancer. 2006 Oct;45(10):950-6.
19. Müller MC, Saglio G, Lin F, Pfeifer H, Press RD, Tubbs RR, Paschka P, Gottardi E, O'Brien SG, Ottmann OG, Stockinger H, Wiecek L, Merx K, König H, Schwindel U, Hehlmann R, Hochhaus A.  
An international study to standardize the detection and quantitation of BCR-ABL transcripts from stabilized peripheral blood preparations by quantitative RT-PCR.  
Haematologica 2007; 92(7):970-973.
20. Müller MC, Erben P, Saglio G, Gottardi E, Nyvold CG, Schenk T, Ernst T, Lauber S, Kruth J, Hehlmann R, Hochhaus A.  
Harmonization of BCR-ABL mRNA quantification using a uniform multifunctional control plasmid in 37 international laboratories.  
Leukemia 2008;22(1):96-102. Epub 2007 Oct 18.
21. Hochhaus A, Kantarjian HM, Baccarani M, Lipton JH, Apperley JF, Druker BJ, Facon T, Goldberg SL, Cervantes F, Niederwieser D, Silver RT, Stone RM, Hughes TP, Müller MC, Ezzeddine R, Countouriotis AM, Shah NP.  
Dasatinib induces notable hematologic and cytogenetic responses in chronic phase chronic myeloid leukemia after failure of imatinib therapy.  
Blood 2007;109(6):2303-9.
22. Fruehauf S, Topaly J, Buss EC, Fischer T, Ottmann OG, Emmerich B, Müller MC, Schuld P, Balleisen L, Hehlmann R, Ho AD, Hochhaus A.

Imatinib combined with mitoxantrone/etoposide and cytarabine is an effective induction therapy for patients with chronic myeloid leukemia in myeloid blast crisis.  
Cancer 2007; 109(8):1543-1549.

23. Hochhaus A, Erben P, Ernst T, Mueller MC.  
Resistance to targeted therapy in chronic myelogenous leukemia.  
Semin Hematol. 2007 Jan;44(1 Suppl 1):15-24.
24. Guilhot F, Apperley J, Kim DW, Bullorsky EO, Baccarani M, Roboz GJ, Amadori S, de Souza CA, Lipton JH, Hochhaus A, Heim D, Larson RA, Branford S, Müller MC, Agarwal P, Gollerkeri A, Talpaz M  
Dasatinib induces significant hematologic and cytogenetic responses in patients with imatinib-resistant or -intolerant chronic myeloid leukemia in accelerated phase.  
Blood 2007; 109(10):4143-50.
25. Fabarius A, Haferlach C, Müller MC, Erben P, Lahaye T, Giehl M, Frank O, Seifarth W, Hehlmann R, Hochhaus A.  
Dynamics of cytogenetic aberrations in Philadelphia chromosome positive and negative hematopoiesis during dasatinib therapy of chronic myeloid leukemia patients after imatinib failure.  
Haematologica 2007; 92(6):834-837.
26. Saldanha J, Silvy M, Beaufils N, Arlinghaus R, Barbany G, Branford S, Cayuela JM, Cazzaniga G, Gonzalez M, Grimwade D, Kairisto V, Miyamura K, Lawler M, Lion T, Macintyre E, Mahon FX, Müller MC, Ostergaard M, Pfeifer H, Saglio G, Sawyers C, Spinelli O, van der Velden V, Wang JQ, Zoi K, Patel V, Phillips P, Matejtschuk P, Gabert J.  
Characterization of a reference material for BCR-ABL (M-BCR) mRNA quantitation by real-time amplification assays: towards new standards for gene expression measurements.  
Leukemia 2007; 21(7):1481-1487.
27. Fabarius A, Giehl M, Frank O, Spiess B, Zheng C, Müller MC, Weiss C, Duesberg P, Hehlmann R, Hochhaus A, Seifarth W.  
Centrosome aberrations after nilotinib and imatinib treatment in vitro are associated with mitotic spindle defects and genetic instability.  
Br J Haematol 2007; 138(3):369-373.
28. Metzgeroth G, Walz C, Score J, Siebert R, Schnittger S, Haferlach C, Popp H, Haferlach T, Erben P, Mix J, Müller MC, Beneke H, Muller L, Del Valle F, Aulitzky WE, Wittkowsky G, Schmitz N,

Schulte C, Muller-Hermelink K, Hodges E, Whittaker SJ, Diecker F, Dohner H, Schuld P, Hehlmann R, Hochhaus A, Cross NC, Reiter A.  
Recurrent finding of the FIP1L1-PDGFR fusion gene in eosinophilia-associated acute myeloid leukemia and lymphoblastic T-cell lymphoma.  
*Leukemia* 2007; 21(6):1183-1188.

29. Kramer A, Reiter A, Kruth J, Erben P, Hochhaus A, Müller M, Cross NC, Jones AV, Ho AD, Hensel M.  
JAK2-V617F mutation in a patient with Philadelphia-chromosome-positive chronic myeloid leukaemia.  
*Lancet Oncol.* 2007 Jul;8(7):658-60.
30. Ernst T, Hoffmann J, Erben P, Hehlmann R, Hochhaus A, Müller MC.  
Suitability of the PAXgene<sup>R</sup> system to stabilize bone marrow RNA in imatinib-resistant patients with chronic myeloid leukemia.  
*Clin Chem Lab Med* 2008;46:318-322.
31. Ernst T, Hoffmann J, Erben P, Hanfstein B, Leitner A, Hehlmann R, Hochhaus A, Müller MC  
ABL single nucleotide polymorphisms may masquerade as BCR-ABL mutations associated with resistance to tyrosine kinase inhibitors in CML patients.  
*Haematologica* 2008;93(6):1389-93.
32. Ernst T, Erben P, Müller MC, Paschka P, Schenk T, Hoffmann J, Kreil S, La Rosée P, Hehlmann R, Hochhaus A.  
Dynamics of BCR-ABL mutated clones prior to hematologic or cytogenetic resistance to imatinib.  
*Haematologica* 2008;93:186-192.
33. Lundán T, Juvonen V, Müller MC, Mustjoki S, Lakkala T, Kairisto V, Hochhaus A, Knuutila S, Porkka K.  
Comparison of bone marrow high mitotic index metaphase fluorescence in situ hybridization to peripheral blood and bone marrow real time quantitative polymerase chain reaction on the International Scale for detecting residual disease in chronic myeloid leukemia.  
*Haematologica* 2008;93:178-85.
34. La Rosée P, Holm-Eriksen S, König H, Härtel N, Ernst T, Debatin J, Müller MC, Erben P, Binckebanck A, Wunderle L, Shou Y, Dugan M, Hehlmann R, Ottmann OG, Hochhaus A.  
Phospho-CRKL monitoring for the assessment of BCR-ABL activity in imatinib-resistant chronic myeloid leukemia or Ph<sup>+</sup> acute lymphoblastic leukemia patients treated with nilotinib.  
*Haematologica* 2008;93:765-9.

35. Hochhaus A, Baccarani M, Deininger M, Apperley JF, Lipton JH, Goldberg SL, Corm S, Shah NP, Cervantes F, Silver RT, Niederwieser D, Stone RM, Dombret H, Larson RA, Roy L, Hughes T, Müller MC, Ezzeddine R, Countouriotis AM, Kantarjian HM.  
Dasatinib induces durable cytogenetic responses in patients with chronic myelogenous leukemia in chronic phase with resistance or intolerance to imatinib.  
Leukemia 2008;22(6):1200-6.
36. Branford S, Fletcher L, Cross NC, Müller MC, Hochhaus A, Kim DW, Radich JP, Saglio G, Pane F, Kamel-Reid S, Wang YL, Press RD, Lynch K, Rudzki Z, Goldman JM, Hughes T.  
Desirable performance characteristics for BCR-ABL measurement on an international reporting scale to allow consistent interpretation of individual patient response and comparison of response rates between clinical trials.  
Blood 2008;112(8):3330-8.
37. Rohrbacher M, Berger U, Hochhaus A, Metzgeroth G, Adam K, Lahaye T, Saussele S, Müller MC, Hasford J, Heimpel H, Hehlmann R.  
Clinical trials underestimate the age of chronic myeloid leukemia (CML) patients. Incidence and median age of Ph/BCR-ABL-positive CML and other chronic myeloproliferative disorders in a representative area in Germany.  
Leukemia. 2009 Mar;23(3):602-4. Epub 2008 Sep 11
38. Erben P, Horisberger K, Muessle B, Müller MC, Treschl A, Ernst T, Kähler G, Ströbel P, Wenz F, Kienle P, Post S, Hochhaus A, Willeke F, Hofheinz RD.  
mRNA expression of platelet-derived growth factor receptor-beta and C-KIT: correlation with pathologic response to cetuximab-based chemoradiotherapy in patients with rectal cancer.  
Int J Radiat Oncol Biol Phys. 2008 Dec 1;72(5):1544-50.
39. Hochhaus A, O'Brien SG, Guilhot F, Druker BJ, Branford S, Foroni L, Goldman JM, Müller MC, Radich JP, Rudoltz M, Mone M, Gathmann I, Hughes TP, Larson RA; IRIS Investigators.  
Six-year follow-up of patients receiving imatinib for the first-line treatment of chronic myeloid leukemia.  
Leukemia. 2009 Jun;23(6):1054-61. Epub 2009 Mar 12
40. Kantarjian H, Cortes J, Kim DW, Dorlhiac-Llacer P, Pasquini R, DiPersio J, Müller MC, Radich JP, Khoury HJ, Khoroshko N, Bradley-Garelik MB, Zhu C, Tallman MS.  
Phase 3 study of dasatinib 140 mg once daily versus 70 mg twice daily in patients with chronic myeloid leukemia in accelerated phase resistant or intolerant to imatinib: 15-month median follow-up.  
Blood. 2009 Jun 18;113(25):6322-9. Epub 2009 Apr 15

41. Ernst T, Gruber FX, Pelz-Ackermann O, Maier J, Pfirrmann M, Müller MC, Mikkola I, Porkka K, Niederwieser D, Hochhaus A, Lange T.  
A co-operative evaluation of different methods of detecting BCR-ABL kinase domain mutations in patients with chronic myeloid leukemia on second-line dasatinib or nilotinib therapy after failure of imatinib.  
Haematologica. 2009 Sep;94(9):1227-35. Epub 2009 Jul 16.
42. Hochhaus A, Müller MC, Radich J, Branford S, Kantarjian HM, Hanfstein B, Rousselot P, Kim DW, Lipton JH, Bleickardt E, Lambert A, Hughes TP.  
Dasatinib-associated major molecular responses in patients with chronic myeloid leukemia in chronic phase following imatinib failure: response dynamics and predictive value.  
Leukemia. 2009 Sep;23(9):1628-33. Epub 2009 Jul 30.
43. Hughes T, Saglio G, Branford S, Soverini S, Kim DW, Müller MC, Martinelli G, Cortes J, Beppu L, Gottardi E, Kim D, Erben P, Shou Y, Haque A, Gallagher N, Radich J, Hochhaus A.  
Impact of baseline BCR-ABL mutations on response to nilotinib in patients with chronic myeloid leukemia in chronic phase.  
J Clin Oncol. 2009 Sep 1;27(25):4204-10. Epub 2009 Aug 3.
44. Müller MC, Cross NC, Erben P, Schenk T, Hanfstein B, Ernst T, Hehlmann R, Branford S, Saglio G, Hochhaus A.  
Harmonization of molecular monitoring of CML therapy in Europe.  
Leukemia. 2009 Nov;23(11):1957-63. Epub 2009 Aug 27
45. Müller MC, Cortes JE, Kim DW, Druker BJ, Erben P, Pasquini R, Branford S, Hughes TP, Radich JP, Ploughman L, Mukhopadhyay J, Hochhaus A.  
Dasatinib treatment of chronic phase chronic myeloid leukemia: analysis of responses according to preexisting BCR-ABL mutations.  
Blood. 2009 Dec 3;114(24):4944-53. Epub 2009 Sep 24
46. Nicolini FE, Mauro MJ, Martinelli G, Kim DW, Soverini S, Müller MC, Hochhaus A, Cortes J, Chuah C, Dufva IH, Apperley JF, Yagasaki F, Pearson JD, Peter S, Sanz Rodriguez C, Preudhomme C, Giles F, Goldman JM, Zhou W.  
Epidemiologic study on survival of chronic myeloid leukemia and Ph(+) acute lymphoblastic leukemia patients with BCR-ABL T315I mutation.  
Blood. 2009 Dec 17;114(26):5271-8. Epub 2009 Oct 20.
47. Hochhaus A, Schenk T, Erben P, Ernst T, La Rosée P, Müller MC.  
Cause and management of therapy resistance.  
Best Pract Res Clin Haematol. 2009 Sep;22(3):367-79.

48. Saussele S, Lauseker M, Gratwohl A, Beelen DW, Bunjes D, Schwerdtfeger R, Kolb HJ, Ho AD, Falge C, Holler E, Schlimok G, Zander AR, Arnold R, Kanz L, Dengler R, Haferlach C, Schlegelberger B, Pfirrmann M, Müller MC, Schnittger S, Leitner A, Pletsch N, Hochhaus A, Hasford J, Hehlmann R.  
Allogeneic hematopoietic stem cell transplantation (alloSCT) for chronic myeloid leukemia in the imatinib era; evaluation of its impact within a subgroup of the randomized German CML Study IV.  
Blood. 2010 Mar 11;115(10):1880-5. Epub 2009 Nov 18.
49. Erben P, Gosenca D, Müller MC, Reinhard J, Score J, Del Valle F, Walz C, Mix J, Metzgeroth G, Ernst T, Haferlach C, Cross NC, Hochhaus A, Reiter A.  
Screening for diverse PDGFRA or PDGFRB by fusion genes is facilitated by generic quantitative RT-PCR.  
Haematologica 2010 May;95(5):738-44. Epub 2010 Jan 27
50. Lilly MB, Ottmann OG, Shah NP, Larson RA, Reiffers JJ, Ehninger G, Müller MC, Charbonnier A, Bullorsky E, Dombret H, Brigid Bradley-Garelik M, Zhu C, Martinelli G.  
Dasatinib 140 mg once daily versus 70 mg twice daily in patients with Ph-positive acute lymphoblastic leukemia who failed imatinib: Results from a phase 3 study.  
Am J Hematol. 2010 Mar;85(3):164-70.
51. Shah NP, Kim DW, Kantarjian H, Rousselot P, Llacer PE, Enrico A, Vela-Ojeda J, Silver RT, Khoury HJ, Müller MC, Lambert A, Matloub Y, Hochhaus A.  
Potent, transient inhibition of BCR-ABL with dasatinib 100 mg daily achieves rapid and durable cytogenetic responses and high transformation-free survival rates in chronic phase chronic myeloid leukemia patients with resistance, suboptimal response or intolerance to imatinib.  
Haematologica. 2010 Feb;95(2):232-40.
52. Burchert A, Müller MC, Kostrewa P, Erben P, Bostel T, Liebler S, Hehlmann R, Neubauer A, Hochhaus A.  
Sustained molecular response with interferon alfa maintenance after induction therapy with imatinib plus interferon alfa in patients with chronic myeloid leukemia.  
J Clin Oncol. 2010 Mar 10;28(8):1429-35. Epub 2010 Feb 8.
53. Giles FJ, Abruzzese E, Rosti G, Kim DW, Bhatia R, Bosly A, Goldberg S, Kam GL, Jagasia M, Mendrek W, Fischer T, Facon T, Dünzinger U, Marin D, Mueller MC, Shou Y, Gallagher NJ, Larson RA, Mahon FX, Baccarani M, Cortes J, Kantarjian HM.  
Nilotinib is active in chronic and accelerated phase chronic myeloid leukemia following failure of imatinib and dasatinib therapy.  
Leukemia. 2010 Jul;24(7):1299-301. Epub 2010 Jun 3.



54. Hughes TP, Hochhaus A, Branford S, Müller MC, Kaeda JS, Foroni L, Druker BJ, Guilhot F, Larson RA, O'Brien SG, Rudoltz MS, Mone M, Wehrle E, Modur V, Goldman JM, Radich JP. Long-term prognostic significance of early molecular response to imatinib in newly diagnosed chronic myeloid leukemia: an analysis from the international randomized study of interferon versus STI571 (IRIS). *Blood*. 2010 Nov 11;116(19):3758-65. Epub 2010 Aug 2.
55. White HE, Matejtschuk P, Rigsby P, Gabert J, Lin F, Wang YL, Branford S, Müller MC, Beaufile N, Beillard E, Colomer D, Dvorakova D, Ehrencrona H, Goh HG, El Housni H, Jones D, Kairisto V, Kamel-Reid S, Kim DW, Langabeer S, Ma ES, Press RD, Romeo G, Wang L, Zoi K, Hughes T, Saglio G, Hochhaus A, Goldman JM, Metcalfe P, Cross NC. Establishment of the 1st World Health Organization International Genetic Reference Panel for quantitation of BCR-ABL mRNA. *Blood*. 2010 Nov 25;116(22):e111-7. Epub 2010 Aug 18.
56. Leitner AA, Hochhaus A, Müller MC. Current Treatment Concepts of CML. *Curr Cancer Drug Targets*. 2011 Jan;11(1):31-43.
57. Hanfstein B, Müller MC, Kreil S, Ernst T, Schenk T, Lorentz C, Schwindel U, Leitner A, Hehlmann R, Hochhaus A. Dynamics of mutant BCR-ABL positive clones after cessation of tyrosine kinase inhibitor therapy. *Haematologica*. 2011 Mar;96(3):360-6. Epub 2010 Dec 6.
58. Hochhaus A, Rosée PL, Müller MC, Ernst T, Cross NC. Impact of BCR-ABL mutations on patients with chronic myeloid leukemia. *Cell Cycle*. 2011 Jan 15;10(2):250-60.
59. Grossmann V, Kohlmann A, Zenger M, Schindela S, Eder C, Weissmann S, Schnittger S, Kern W, Müller MC, Hochhaus A, Haferlach T, Haferlach C. A deep-sequencing study of chronic myeloid leukemia patients in blast crisis (BC-CML) detects mutations in 76.9% of cases. *Leukemia*. 2011 Mar;25(3):557-60. Epub 2011 Jan 28
60. Hehlmann R, Lauseker M, Jung-Munkwitz S, Leitner A, Müller MC, Pletsch N, Proetel U, Haferlach C, Schlegelberger B, Balleisen L, Hänel M, Pfirrmann M, Krause SW, Nerl C, Pralle H, Gratwohl A, Hossfeld DK, Hasford J, Hochhaus A, Saübele S. Tolerability-Adapted Imatinib 800 mg/d Versus 400 mg/d Versus 400 mg/d Plus Interferon- $\alpha$  in Newly Diagnosed Chronic Myeloid Leukemia. *J Clin Oncol*. 2011 Apr 20;29(12):1634-1642. Epub 2011 Mar 21.

61. Soverini S, Hochhaus A, Nicolini FE, Gruber F, Lange T, Saglio G, Pane F, Müller MC, Ernst T, Rosti G, Porkka K, Baccarani M, Cross NC, Martinelli G.  
Bcr-Abl kinase domain mutation analysis in chronic myeloid leukemia patients treated with tyrosine kinase inhibitors: recommendations from an expert panel on behalf of European LeukemiaNet.  
Blood. 2011 Aug 4;118(5):1208-15. Epub May 11.
62. How GF, Müller MC, Lim LC.  
Acquisition of the novel ABL kinase domain mutation T315L in a relapsed Philadelphia-positive acute lymphoblastic leukemia patient.  
Leuk Res. 2012 Jan;36(1):e20-1. Epub 2011 Sep 6
63. Hurtz C, Hatzi K, Cerchietti L, Braig M, Park E, Kim YM, Herzog S, Ramezani-Rad P, Jumaa H, Müller MC, Hofmann WK, Hochhaus A, Ye BH, Agarwal A, Druker BJ, Shah NP, Melnick AM, Müschen M.  
BCL6-mediated repression of p53 is critical for leukemia stem cell survival in chronic myeloid leukemia.  
J Exp Med. 2011 Oct 24;208(11):2163-74. Epub 2011 Sep 12.
64. Nicolini FE, Basak GW, Soverini S, Martinelli G, Mauro MJ, Müller MC, Hochhaus A, Chuah C, Dufva IH, Rege-Cambrin G, Saglio G, Michallet M, Labussière H, Morisset S, Hayette S, Etienne G, Olavarria E, Zhou W, Peter S, Apperley JF, Cortes J.  
Allogeneic stem cell transplantation for patients harboring T315I BCR-ABL mutated leukemias.  
Blood. 2011 Nov 17;118(20):5697-700. Epub 2011 Sep 16.
65. Fabarius A, Leitner A, Hochhaus A, Müller MC, Hanfstein B, Haferlach C, Göhring G, Schlegelberger B, Jotterand M, Reiter A, Jung-Munkwitz S, Proetel U, Schwaab J, Hofmann WK, Schubert J, Einsele H, Ho AD, Falge C, Kanz L, Neubauer A, Kneba M, Stegelmann F, Pfreundschuh M, Waller CF, Spiekermann K, Baerlocher GM, Lauseker M, Pfirrmann M, Hasford J, Saussele S, Hehlmann R; for the Schweizerische Arbeitsgemeinschaft für Klinische Krebsforschung (SAKK) and the German CML Study Group.  
Impact of additional cytogenetic aberrations at diagnosis on prognosis of CML: long-term observation of 1151 patients from the randomized CML Study IV.  
Blood. 2011 Dec 22;118(26):6760-6768. Epub 2011 Oct 28.
66. Ernst T, La Rosée P, Müller MC, Hochhaus A.  
BCR-ABL mutations in chronic myeloid leukemia.  
Hematol Oncol Clin North Am. 2011 Oct;25(5):997-1008, v-vi.

67. Härtel N, Klag T, Hanfstein B, Mueller MC, Schenk T, Erben P, Hochhaus A, La Rosée P. Enhanced ABL-inhibitor-induced MAPK-activation in T315I-BCR-ABL-expressing cells: a potential mechanism of altered leukemogenicity. *J Cancer Res Clin Oncol*. 2012 Feb;138(2):203-12. Epub 2011 Nov 17
68. Early molecular and cytogenetic response is predictive for long-term progression-free and overall survival in chronic myeloid leukemia (CML). Hanfstein B, Müller MC, Hehlmann R, Erben P, Lauseker M, Fabarius A, Schnittger S, Haferlach C, Göhring G, Proetel U, Kolb HJ, Krause SW, Hofmann WK, Schubert J, Einsele H, Dengler J, Hänel M, Falge C, Kanz L, Neubauer A, Kneba M, Stegelmann F, Pfreundschuh M, Waller CF, Branford S, Hughes TP, Spiekermann K, Baerlocher GM, Pffirmann M, Hasford J, Saußebe S, Hochhaus A. *Leukemia*. 2012 Sep; 26(9):2096-102. Epub 2012 Mar 26.
69. SNP array analysis of acute promyelocytic leukemia may be of prognostic relevance and identifies a potential high risk group with recurrent deletions on chromosomal subband 1q31.3. Nowak D, Klaumuenzer M, Hanfstein B, Mossner M, Nolte F, Nowak V, Oblaender J, Hecht A, Hütter G, Ogawa S, Kohlmann A, Haferlach C, Schlegelberger B, Braess J, Seifarth W, Fabarius A, Erben P, Saussele S, Müller MC, Reiter A, Buechner T, Weiss C, Hofmann WK, Lengfelder E. *Genes Chromosomes Cancer*. 2012 Aug;51(8):756-67. Epub 2012 Apr 9.
70. Standardized definitions of molecular response in chronic myeloid leukemia. Cross NC, White HE, Müller MC, Saglio G, Hochhaus A. *Leukemia*. 2012 Oct;26(19):2172-5. Epub 2012 Apr 16.
71. The quantitative level of T315I mutated BCR-ABL predicts for major molecular response to second line nilotinib or dasatinib treatment in patients with chronic myeloid leukemia. Lange T, Ernst T, Gruber F, Maier J, Cross M, Müller MC, Niederwieser D, Hochhaus A, Pffirmann M. *Haematologica*. 2013 May;98(5):714-7. Epub 2012 Oct 12.
72. Chronic myeloid leukemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Baccarani M, Pileri S, Steegmann JL, Muller M, Soverini S, Dreyling M; ESMO Guidelines Working Group. *Ann Oncol*. 2012 Oct;23 Suppl 7:vii72-7. No abstract available.

73. Initial molecular response at 3 months may predict both response and event-free survival at 24 months in imatinib-resistant or -intolerant patients with Philadelphia chromosome-positive chronic myeloid leukemia in chronic phase treated with nilotinib.  
Branford S, Kim DW, Soverini S, Haque A, Shou Y, Woodman RC, Kantarjian HM, Martinelli G, Radich JP, Saglio G, Hochhaus A, Hughes TP, Müller MC.  
J Clin Oncol. 2012;30(35):4323-9. Epub 2012 Oct 29.
74. Model-based decision rules reduce the risk of molecular relapse after cessation of tyrosine kinase inhibitor therapy in chronic myeloid leukemia.  
Horn M, Glauche I, Müller MC, Hehlmann R, Hochhaus A, Loeffler M, Roeder I.  
Blood. 2013;121(2):378-84. Epub 2012 Nov 21.
75. The hOCT1 SNPs M420del and M408V alter imatinib uptake and M420del modifies clinical outcome in imatinib-treated chronic myeloid leukaemia.  
Giannoudis A, Wang L, Jorgensen AL, Xinarianos G, Davies A, Pushpakom S, Liloglou T, Zhang JE, Austin G, Holyoake TL, Foroni L, Kottaridis PD, Müller MC, Pirmohamed M, Clark RE.  
Blood. 2013;121(4):628-37. Epub 2012 Dec 5.
76. Interpretation of molecular diagnostics results in routine monitoring for chronic myeloid leukemia: recommendations from a German expert panel.  
Hochhaus A, Müller MC, Lange T, Thiede C, Haferlach T.  
Laboratoriumsmedizin-Journal of Laboratory Medicine. 2012;36(6):405-11.
77. Establishment and validation of analytical reference panels for the standardization of quantitative BCR-ABL1 measurements on the international scale.  
White HE, Hedges J, Bendit I, Branford S, Colomer D, Hochhaus A, Hughes T, Kamel-Reid S, Kim DW, Modur V, Müller MC, Pagnano KB, Pane F, Radich J, Cross NC, Labourier E.  
Clin Chem. 2013;59(6):938-48. Epub 2013 Mar 7.
78. Genomic instability may originate from imatinib-refractory chronic myeloid leukemia stem cells.  
Bolton-Gillespie E, Schemionek M, Klein HU, Flis S, Hoser G, Lange T, Nieborowska-Skorska M, Maier J, Kerstiens L, Koptyra M, Müller MC, Modi H, Stoklosa T, Seferynska I, Bhatia R, Holyoake TL, Koschmieder S, Skorski T.  
Blood. 2013;121(20):4175-83. Epub 2013 Mar 29.
79. Rapid initial decline in BCR-ABL1 is associated with superior responses to second-line nilotinib in patients with chronic-phase chronic myeloid leukemia.  
Stein AM, Martinelli G, Hughes TP, Müller MC, Beppu L, Gottardi E, Branford S, Soverini S, Woodman RC, Hochhaus A, Kim DW, Saglio G, Radich JP.  
BMC Cancer. 2013 Apr 2;13(1):173. [Epub ahead of print]

80. Improved tolerability by a modified intermittent treatment schedule of dasatinib for patients with chronic myeloid leukemia resistant or intolerant to imatinib.  
La Rosée P, Martiat P, Leitner A, Klag T, Müller MC, Erben P, Schenk T, Saussele S, Hochhaus A.  
Ann Hematol. 2013 Oct;92(10):1345-50. Epub 2013 Apr 28.
81. The BCR-ABL T315I mutation compromises survival in chronic phase chronic myelogenous leukemia patients resistant to tyrosine kinase inhibitors, in a matched pair analysis.  
Nicolini FE, Ibrahim AR, Soverini S, Martinelli G, Müller MC, Hochhaus A, Dufva IH, Kim DW, Cortes J, Mauro MJ, Chuah C, Labussière H, Morisset S, Roche-Lestienne C, Lippert E, Hayette S, Peter S, Zhou W, Maguer-Satta V, Michallet M, Goldman J, Apperley JF, Mahon FO, Marin D, Etienne G.  
Haematologica. 2013 98(10):1510-6. Epub 2013 May 28.
82. European LeukemiaNet recommendations for the management of chronic myeloid leukemia: 2013.  
Baccarani M, Deininger MW, Rosti G, Hochhaus A, Soverini S, Apperley JF, Cervantes F, Clark RE, Cortes JE, Guilhot F, Hjorth-Hansen H, Hughes TP, Kantarjian HM, Kim DW, Larson RA, Lipton JH, Mahon FX, Martinelli G, Mayer J, Müller MC, Niederwieser D, Pane F, Radich JP, Rousselot P, Saglio G, Saúbele S, Schiffer C, Silver R, Simonsson B, Steegmann JL, Goldman JM, Hehlmann R.  
Blood. 2013;122(6):872-84. Epub 2013 Jun 26.
83. The ne plus ultra for deep BCR-ABL sequencing?  
Müller MC.  
Blood. 2013 Aug 29;122(9):1536-7.
84. Implications of BCR-ABL1 kinase domain-mediated resistance in chronic myeloid leukemia.  
Soverini S, Branford S, Nicolini FE, Talpaz M, Deininger MW, Martinelli G, Müller MC, Radich JP, Shah NP.  
Leuk Res. 2014;38(1):10-20. Epub 2013 Sep 23
85. Younger patients with chronic myeloid leukemia do well in spite of poor prognostic indicators: results from the randomized CML study IV.  
Kalmanti L, Saussele S, Lauseker M, Proetel U, Müller MC, Hanfstein B, Schreiber A, Fabarius A, Pfirrmann M, Schnittger S, Dengler J, Falge C, Kanz L, Neubauer A, Stegelmann F, Pfreundschuh M, Waller CF, Spiekermann K, Krause SW, Heim D, Nerl C, Hossfeld DK, Kolb HJ, Hochhaus A, Hasford J, Hehlmann R; German Chronic Myeloid Leukemia Study Group; Schweizerische Arbeitsgemeinschaft für Klinische Krebsforschung (SAKK).  
Ann Hematol. 2014;93(1):71-80. Epub 2013 Oct 27

86. Fatal progressive cerebral ischemia in CML under third-line treatment with ponatinib.  
Mayer K, Gielen GH, Willinek W, Müller MC, Wolf D.  
Leukemia. 2014;28(4):976-7. Epub 2013 Oct 30
87. A phase 2 trial of ponatinib in Philadelphia chromosome-positive leukemias.  
Cortes JE, Kim DW, Pinilla-Ibarz J, le Coutre P, Paquette R, Chuah C, Nicolini FE, Apperley JF, Khoury HJ, Talpaz M, DiPersio J, DeAngelo DJ, Abruzzese E, Rea D, Baccarani M, Müller MC, Gambacorti-Passerini C, Wong S, Lustgarten S, Rivera VM, Clackson T, Turner CD, Haluska FG, Guilhot F, Deininger MW, Hochhaus A, Hughes T, Goldman JM, Shah NP, Kantarjian H; PACE Investigators.  
N Engl J Med. 2013;369(19):1783-96. Epub 2013 Nov 1
88. The KIT D816V expressed allele burden for diagnosis and disease monitoring of systemic mastocytosis.  
Erben P, Schwaab J, Metzgeroth G, Horny HP, Jawhar M, Sotlar K, Fabarius A, Teichmann M, Schneider S, Ernst T, Müller MC, Giehl M, Marx A, Hartmann K, Hochhaus A, Hofmann WK, Cross NC, Reiter A.  
Ann Hematol. 2014;93(1):81-8. Epub 2013 Nov 27
89. Deep Molecular Response Is Reached by the Majority of Patients Treated With Imatinib, Predicts Survival, and Is Achieved More Quickly by Optimized High-Dose Imatinib: Results From the Randomized CML-Study IV.  
Hehlmann R, Müller MC, Lauseker M, Hanfstein B, Fabarius A, Schreiber A, Proetel U, Pletsch N, Pfirrmann M, Haferlach C, Schnittger S, Einsele H, Dengler J, Falge C, Kanz L, Neubauer A, Kneba M, Stegelmann F, Pfreundschuh M, Waller CF, Spiekermann K, Baerlocher GM, Ehninger G, Heim D, Heimpel H, Nerl C, Krause SW, Hossfeld DK, Kolb HJ, Hasford J, Sauße S, Hochhaus A.  
J Clin Oncol. 2014;32(5):415-23. Epub 2013 Dec 2
90. Expression of transketolase-like gene 1 (TKTL1) depends on disease phase in patients with chronic myeloid leukaemia (CML).  
Philipp M, Schwaab J, Dietz CT, Hanfstein B, Kalmanti L, Munjal U, Mossner M, Nowak D, Seifarth W, Hofmann WK, Hochhaus A, Müller MC, Erben P.  
J Cancer Res Clin Oncol. 2014;140(3):411-7. Epub 2014 Jan 5
91. MDR1 expression predicts outcome of Ph+ chronic phase CML patients on second-line nilotinib therapy after imatinib failure.  
Agrawal M, Hanfstein B, Erben P, Wolf D, Ernst T, Fabarius A, Saussele S, Purkayastha D, Woodman RC, Hofmann WK, Hehlmann R, Hochhaus A, Müller MC.  
Leukemia. 2014;28(7):1487-85. Epub 2014 Jan 10

92. Older patients with chronic myeloid leukemia ( $\geq 65$  years) profit more from higher imatinib doses than younger patients: a subanalysis of the randomized CML-Study IV.  
Proetel U, Pletsch N, Lauseker M, Müller MC, Hanfstein B, Krause SW, Kalmanti L, Schreiber A, Heim D, Baerlocher GM, Hofmann WK, Lange E, Einsele H, Wernli M, Kremers S, Schlag R, Müller L, Hänel M, Link H, Hertenstein B, Pfirrmann M, Hochhaus A, Hasford J, Hehlmann R, Sauße S; for the German Chronic Myeloid Leukemia Study Group, and the Schweizerische Arbeitsgemeinschaft für Klinische Krebsforschung (SAKK)..  
Ann Hematol. 2014;93(7):1167-76. Epub 2014 Mar 22
93. Velocity of early BCR-ABL transcript elimination as an optimized predictor of outcome in chronic myeloid leukemia (CML) patients in chronic phase on treatment with imatinib.  
Hanfstein B, Shlyakhto V, Lauseker M, Hehlmann R, Saussele S, Dietz C, Erben P, Fabarius A, Proetel U, Schnittger S, Krause SW, Schubert J, Einsele H, Hänel M, Dengler J, Falge C, Kanz L, Neubauer A, Kneba M, Stegelmann F, Pfreundschuh M, Waller CF, Spiekermann K, Baerlocher GM, Pfirrmann M, Hasford J, Hofmann WK, Hochhaus A, Müller MC.  
Leukemia. 2014;28(10):1988-92. Epub 2014 May 6
94. Distinct characteristics of e13a2 versus e14a2 BCR-ABL1 driven chronic myeloid leukemia under first-line therapy with imatinib.  
Hanfstein B, Lauseker M, Hehlmann R, Saussele S, Erben P, Dietz C, Fabarius A, Proetel U, Schnittger S, Haferlach C, Krause SW, Schubert J, Einsele H, Hänel M, Dengler J, Falge C, Kanz L, Neubauer A, Kneba M, Stegelmann F, Pfreundschuh M, Waller CF, Spiekermann K, Baerlocher GM, Pfirrmann M, Hasford J, Hofmann WK, Hochhaus A, Müller MC.  
Haematologica. 2014;99(9):1441-7. Epub 2014 May 16
95. Equivalence of BCR-ABL transcript levels with complete cytogenetic remission in patients with chronic myeloid leukemia in chronic phase.  
Lauseker M, Hanfstein B, Haferlach C, Schnittger S, Pfirrmann M, Fabarius A, Schlegelberger B, Sauße S, Dietz CT, Erben P, Hehlmann R, Hasford J, Hochhaus A, Müller MC.  
J Cancer Res Clin Oncol. 2014;140(11):1965-9. Epub 2014 Jun 22
96. A certified plasmid reference material for the standardisation of BCR-ABL1 mRNA quantification by real-time quantitative PCR.  
White H, Deprez L, Corbisier P, Hall V, Lin F, Mazoua S, Trapmann S, Aggerholm A, Andrikovics H, Akiki S, Barbany G, Boeckx N, Bench A, Catherwood M, Cayuela JM, Chudleigh S, Clench T, Colomer D, Daraio F, Dulucq S, Farrugia J, Fletcher L, Foroni L, Ganderton R, Gerrard G, Gineikienė E, Hayette S, El Housni H, Izzo B, Jansson M, Johnels P, Jurcek T, Kairisto V, Kizilors A, Kim DW, Lange T, Lion T, Polakova KM, Martinelli G, McCarron S, Merle PA, Milner B, Mitterbauer-Hohendanner G, Nagar M, Nickless G, Nomdedéu J, Nymoen DA, Leibundgut EO, Ozbek U, Pajič T, Pfeifer H, Preudhomme C, Raudsepp K, Romeo G, Sacha T, Talmaci R, Touloumenidou T, Van der Velden VH, Waits P, Wang L, Wilkinson E, Wilson G,

Wren D, Zadro R, Ziermann J, Zoi K, Müller MC, Hochhaus A, Schimmel H, Cross NC, Emons H.

Leukemia. 2014 Jul 18. [Epub ahead of print]

97. Dual Glutathione-S-Transferase- $\theta$ 1 and - $\mu$ 1 Gene Deletions Determine Imatinib Failure in Chronic Myeloid Leukemia.

Davies A, Giannoudis A, Zhang JE, Austin G, Wang L, Holyoake TL, Müller MC, Foroni L, Kottaridis PD, Pirmohamed M, Clark RE.

Clin Pharmacol Ther. 2014;96(6):694-703. Epub 2014 Sep 4